

Non-Perturbative QCD at Finite Temperature

Pok Man Lo

University of Pittsburgh

Jlab Hugs student talk, 6-20-2008

personal information

- email: pol4@pitt.edu
- institution: University of Pittsburgh
- advisor: Eric Swanson

outline

1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory

- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

Outline

1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory

- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

Introduction

The progress of theoretical physics

Introduction

The progress of theoretical physics

The search for exact solution:

Introduction

The progress of theoretical physics

The search for exact solution:

- Newtonian physics: 3-body problem was insoluble

Introduction

The progress of theoretical physics

The search for exact solution:

- Newtonian physics: 3-body problem was insoluble
- QED: 2-body and 1-body problem was insoluble

Introduction

The progress of theoretical physics

The search for exact solution:

- Newtonian physics: 3-body problem was insoluble
- QED: 2-body and 1-body problem was insoluble
- now: 0-body, namely, the vacuum is insoluble

Introduction

The progress of theoretical physics


The search for exact solution:

- Newtonian physics: 3-body problem was insoluble
- QED: 2-body and 1-body problem was insoluble
- now: 0-body, namely, the vacuum is insoluble

No body is too many!!


Introduction

Difficult in calculating arbitrarily complicated diagram



Introduction

Difficult in calculating arbitrarily complicated diagram



Impossible to **sum** all the diagrams

Introduction

Perturbation: expansion in α

Introduction

Perturbation: expansion in α

- is not useful if α is not small

Introduction

Perturbation: expansion in α

- is not useful if α is not small
- cannot explain non-perturbative phenomena:
e.g bound state formation and spontaneous symmetry breaking

Introduction

Perturbation: expansion in α

- is not useful if α is not small
- cannot explain non-perturbative phenomena:
e.g bound state formation and spontaneous symmetry breaking

classical mechanics example: mass attached to a spring: $A \sin(\sqrt{\frac{k}{m}} t)$

Outline


1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory


- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

illustration of the failure of perturbation



...an innocent looking function


illustration of the failure of perturbation



...an innocent looking function
using Taylor expansion, we write

$$f(x) = A_0 + A_1x + A_2x^2 + \dots$$

illustration of the failure of perturbation



...an innocent looking function
using Taylor expansion, we write

$$f(x) = A_0 + A_1x + A_2x^2 + \dots$$

A_0, A_1, A_2, \dots are all strictly 0!

illustration of the failure of perturbation

differential equation...

illustration of the failure of perturbation

differential equation...

$$f(x) + x^3 f' = 0$$

illustration of the failure of perturbation

differential equation...

$$f(x) + x^3 f' = 0$$

$$f(x) = \exp^{-\frac{1}{x^2}}$$

illustration of the failure of perturbation

differential equation...

$$f(x) + x^3 f' = 0$$

$$f(x) = \exp^{-\frac{1}{x^2}}$$

any other method will work, other than perturbation!

Different Philosophy

Perturbative Vs Non-perturbative:

Different Philosophy

Perturbative Vs Non-perturbative:

perturbative method: sum **all** diagrams up to **certain order** in α

Different Philosophy

Perturbative Vs Non-perturbative:

perturbative method: sum **all** diagrams up to **certain order** in α

non-perturbative method: sum a **certain class** of diagram to **all** order

Different Philosophy

Perturbative Vs Non-perturbative:

perturbative method: sum **all** diagrams up to **certain order** in α

non-perturbative method: sum a **certain class** of diagram to **all** order

in some sense, non-perturbative method includes the whole perturbation method as the latter is just a classification of diagram by the coupling constant α

Different Philosophy

Perturbative Vs Non-perturbative:

perturbative method: sum **all** diagrams up to **certain order** in α

non-perturbative method: sum a **certain class** of diagram to **all** order

in some sense, non-perturbative method includes the whole perturbation method as the latter is just a classification of diagram by the coupling constant α

approximation \neq perturbation!

Different Philosophy

Perturbative Vs Non-perturbative:

perturbative method: sum **all** diagrams up to **certain order** in α

non-perturbative method: sum a **certain class** of diagram to **all** order

in some sense, non-perturbative method includes the whole perturbation method as the latter is just a classification of diagram by the coupling constant α

approximation \neq perturbation!

QFT contains more than just the S-matrix and perturbation!

Outline

1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory

- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

motivation

perturbation does not help as we want to sum diagrams to all order in α

new tools other than perturbation!

in some sense, we need exact relations among diagrams!

two particularly useful ones:

- Diagrammatics
- Schwinger Dyson Equation

method of partial sum

summing a particular class of diagrams to all order


method of partial sum

summing a particular class of diagrams to all order

to be explicit, let's consider the diagrams for constructing a propagator

diagrammatics of propagator

in general, to construct the propagator, we need to sum...




diagrammatics of propagator

in general, we cannot sum them all!

diagrammatics of propagator

in general, we cannot sum them all!

but let's focus on the following class of diagram:




$$= \frac{|}{1 - \begin{array}{c} \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \end{array}}$$

diagrammatics of propagator

in general, we cannot sum them all!

but let's focus on the following class of diagram:



$$\begin{aligned}
 &= \frac{\text{Diagram 1}}{1 - \text{Diagram 3}} \\
 &= \frac{1}{\text{Diagram 3}} \cdot \text{Diagram 3}
 \end{aligned}$$

diagrammatics of propagator

the above procedure should be compared with the summing of geometric series:

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

diagrammatics of propagator

Two comments:

diagrammatics of propagator

Two comments:

inspired by the series, we know we can do better:


diagrammatics of propagator

Two comments:


inspired by the series, we know we can do better:

$$\begin{aligned}
 & \text{Diagram 1} = \text{Diagram 2} + \text{Diagram 3} + \text{Diagram 4} + \text{Diagram 5} + \dots \\
 & = \frac{1}{\text{Diagram 6} - (\text{Diagram 7} + \text{Diagram 8})}
 \end{aligned}$$

diagrammatics of propagator




diagrammatics of propagator




$$\Longleftrightarrow \not{p} - m$$

diagrammatics of propagator



diagrammatics of propagator



\Longleftrightarrow dynamical mass

Schwinger Dyson Equation

an exact relations among n-point functions

Schwinger Dyson Equation

an exact relations among n-point functions

basic idea:

$$\int dx \frac{d}{dx} f(x) = 0$$

Schwinger Dyson Equation

an exact relations among n-point functions

basic idea:

$$\int dx \frac{d}{dx} f(x) = 0$$

$$\mathcal{Z} = \int D\psi D\bar{\psi} DGe^{i(S + \int \bar{\eta}\psi + \bar{\psi}\eta + \dots)} / \int D\psi D\bar{\psi} DGe^{i(S)}$$

Schwinger Dyson Equation

an exact relations among n-point functions

basic idea:

$$\int dx \frac{d}{dx} f(x) = 0$$

$$\mathcal{Z} = \int D\psi D\bar{\psi} DGe^{i(S + \int \bar{\eta}\psi + \bar{\psi}\eta + \dots)} / \int D\psi D\bar{\psi} DGe^{i(S)}$$

using the fact

$$\int D\psi D\bar{\psi} DG \left(\frac{\delta}{\delta \bar{\psi}} e^{i(S + \int \bar{\eta}\psi + \bar{\psi}\eta + \dots)} \right) = 0$$
$$\mathcal{Z} = e^{i\mathcal{W}}$$

Schwinger Dyson Equation

as an illustration, for the Hamiltonian:

$$H = \int d^3x \psi_{\vec{x}}^\dagger \gamma^0 [-i\vec{\gamma} \cdot \nabla + m] \psi_{\vec{x}} - G \int d^3x d^3y V_{\vec{x}\vec{y}} \psi_{\vec{x}}^\dagger T^a \psi_{\vec{x}} \psi_{\vec{y}}^\dagger T^a \psi_{\vec{y}}$$

Schwinger Dyson Equation

$$(i\gamma \cdot \partial_x - m) \frac{\delta^2 \mathcal{W}}{\delta \bar{\eta}_x \delta \eta_y} + 2G \int d^4z V \gamma^0 T^a i \frac{\delta^2 \mathcal{W}}{\delta \bar{\eta}_x \delta \eta_z} \gamma^0 T^a \frac{\delta^2 \mathcal{W}}{\delta \bar{\eta}_z \delta \eta_y} = \delta_{xy}$$

Outline

1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory

- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

Gap Equation Revisited

to illustrate the main idea above, we look at the gap equation

Gap Equation Revisited

to illustrate the main idea above, we look at the gap equation

consider the Hamiltonian:

$$H = \int d^3x \psi_{\vec{x}}^\dagger \gamma^0 [-i\vec{\gamma} \cdot \nabla + m] \psi_{\vec{x}} - G \int d^3x d^3y V_{\vec{x}\vec{y}} \psi_{\vec{x}}^\dagger T^a \psi_{\vec{x}} \psi_{\vec{y}}^\dagger T^a \psi_{\vec{y}}$$

for the contact case: $V \rightarrow \delta^{(3)}$

this corresponds to the case where the quark exchange an instantaneous gluon locally

Gap Equation Revisited


the gap equation tell you how the quark is dressed according to the Hamiltonian

namely, the dynamical mass generated:

before showing you the answer, we need to perform our final dressing...

the final dressing

in fact we miss some diagram...



the final dressing

The diagram shows an equation for the final dressing of a quark line. On the left is a single quark line with a black dot representing a vertex. This is equal to a sum of two terms, each representing a quark line with a gluon loop (a circle with two wavy lines). The first term has a horizontal line above the loop and a vertical line to the left of the loop, with a minus sign and a '1' above the vertical line. The second term has a horizontal line above the loop and a vertical line to the right of the loop, with a plus sign and a '1' above the vertical line.

the final dressing

The diagram shows an equation for the final dressing of a fermion line. On the left is a single fermion line with a black dot. This is equal to a sum of two terms, each enclosed in a horizontal line labeled '1' above it. The first term is a fermion line with a black dot, followed by a gluon loop (represented by a wavy line) with a black dot on the right side. The second term is a fermion line with a black dot, followed by a gluon loop (represented by a wavy line) with a black dot on the left side. The two terms are separated by a plus sign.

Gap Equation Revisited

the gap equation for the general case:

Generalized Gap Equation


$$M(\vec{k}) = m - G \frac{\text{Tr}[TT]}{N_c} \int \frac{d^3 k'}{(2\pi)^3} V_{\vec{k}' - \vec{k}} \left[\frac{M(\vec{k}')}{E_{\vec{k}'}} - \hat{k}' \cdot \hat{k} \frac{|\vec{k}'| M(\vec{k})}{E_{\vec{k}'} |\vec{k}|} \right]$$

$M(\vec{k})$ in general is a function of \vec{k}

it dictates how the mass is dynamically generated

Gap Equation Revisited

for the contact case, we have...



Outline

1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory

- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

motivation

FTFT is needed to study the physics of QGP, deconfinement and chiral restoration

when calculating observables in QFT, we only calculate the vacuum expectation value

at finite temperature, excited states start to contribute, the interesting quantity should be the **thermal average** of the observables

we expect n_{E_k} to enter QFT

partition function dictates the equilibrium Finite Temperature QFT

Outline

1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory

- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

basics of FTFT

the partition function:

$$\mathcal{Z} = \text{Tr}[e^{-\beta H}] = \int dq \langle q | e^{-\beta H} | q \rangle$$

observables are given by

$$\mathcal{O} = \text{Tr}[e^{-\beta H} \mathcal{O}] = \langle\langle q | \mathcal{O} | q \rangle\rangle$$

working in imaginary time $\tau = it$

a corresponding path integral representation of the partition function, with Periodic/Antiperiodic boundary condition

the boundary condition motivates the use of Matsubara Green's function:

$$\int dk^0 \rightarrow \frac{1}{\beta} \sum_{\omega_n}$$

Outline

1 Non-perturbative QCD at zero temperature

- Introduction to non-perturbative physics
- Tools for studying non-perturbative QCD
 - Diagrammatics
 - Schwinger Dyson Equation
- Illustration: Gap Equation Revisited

2 Finite Temperature Field Theory

- Survey of basic formalism of Finite Temperature of Field Theory
- Finite Temperature Gap Equation

Gap equation in Finite Temperature

Generalized Gap Equation

$$M(\vec{k}) = m - G \frac{\text{Tr}[TT]}{N_c} \int \frac{d^3 k'}{(2\pi)^3} V_{\vec{k}' - \vec{k}} \left[\frac{M(\vec{k}')}{E_{\vec{k}'}} - \hat{k}' \cdot \hat{k} \frac{|\vec{k}'| M(\vec{k})}{E_{\vec{k}'} |\vec{k}|} \right] (1 - 2n_{E_{\vec{k}'}})$$

$$\text{with } n_{E_{\vec{k}}} = \frac{1}{e^{\beta E_{\vec{k}}} + 1}$$

summary

- non-perturbative physics with $e^{\frac{-1}{x^2}}$
- summation of a class of diagram with $\frac{1}{1-x}$
- Schwinger Dyson Equation with $\int dx \frac{d}{dx} f(x)$
- Finite Temperature Field Theory with $Tr[e^{-\beta H}]$

thank you